

CONSENT FOR SILVER DIAMINE FLUORIDE

Facts for consideration:

- Silver Diamine Fluoride (SDF) is an antibacterial liquid used as a treatment option for tooth decay and tooth sensitivity. SDF application may require repeated application.
- **DO NOT EAT 2-3 HOURS PRIOR TO APPOINTMENT. Normal eating or drinking can resume 30 minutes after application.**
- **SDF does not eliminate the need for dental fillings or crowns to repair function or esthetics. Additional procedures will incur a separate fee.**
- I should not be treated with SDF if: 1) I am **allergic to silver** 2) there are painful sores or raw areas on my gums (i.e., ulcerative gingivitis) or anywhere in my mouth (i.e., stomatitis).

Benefits of receiving SDF:

- SDF can help slow the progression of tooth decay and relieve tooth sensitivity.
- SDF can help buy time for those patients who are very young, fearful, or have special needs that may otherwise require sedation for traditional dental treatment.



Risks related to SDF include, but are not limited to:

- **The affected area will stain black permanently.** Healthy tooth structure may stain. Stained tooth structure can be replaced with a filling or a crown.
- **Tooth-colored fillings and crowns may discolor if SDF is applied to them.** Color changes on the surface can normally be polished off. The edge between a tooth and filling may keep the color.
- **If the material comes in contact with the face, skin, lips, or gums, a brown, dark blue, or black stain may appear. This stain causes no harm, but may appear to look like bruising. The stain cannot be washed off and will disappear in 1-3 weeks.**
- If tooth decay is not arrested, the decay will progress. In that case the tooth will require further treatment, such as repeat SDF, a filling or crown, root canal treatment, or extraction.
- You may notice a metallic taste. These side effects may not include all of the possible situations reported by the manufacturer. If you notice other effects, please contact our office.
- **Every reasonable effort will be made to ensure the success of SDF treatment. There is a risk that the procedure will not stop the decay and no guarantee of success is granted or implied.**

Alternatives to SDF, not limited to the following:

- No treatment, which may lead to continued deterioration of tooth structures and cosmetic appearance. Symptoms may increase in severity.
- Depending on the location and extent of the tooth decay, other treatment may include placement of fluoride varnish, a filling or crown, extraction, or referral for advanced treatment modalities.

**I CERTIFY THAT I HAVE READ AND FULLY UNDERSTAND THIS DOCUMENT,
AND ALL MY QUESTIONS WERE ANSWERED:**

Parent Signature: _____

Date: _____