



Garth LeCheminant, M.D. **David Morrow, M.D., F.A.C.S.** **Iuliana Bobanga, M.D. F.A.C.S.** **Jennifer Lemert, N.P.**
2490 S. Woodworth Loop, Ste 450 Palmer, AK 99645 • Phone: (907) 745-8100 Fax: (907) 746-2655 • matsusurgical.com

PREPARATION INSTRUCTIONS FOR COLONOSCOPY

FACILITY INSTRUCTIONS

The facility does not schedule the times for the colonoscopies until the day before.

Someone from the facility will call you **the afternoon before** the colonoscopy with the time to arrive.

____ **Mat-Su Regional Medical Center (907) 861-6858**

____ **Surgery Center of Wasilla (907) 631-3578**

YOU MUST MAKE ARRANGEMENTS FOR SOMEONE YOU KNOW TO DRIVE YOU HOME. You may not drive for the rest of the day once the procedure is completed. **You CANNOT take a cab home.**

MoviPrep ONE DAY Method - COLON CLEANSE INSTRUCTIONS

PLEASE READ THESE INSTRUCTIONS SEVERAL DAYS BEFORE YOUR PROCEDURE.

Purchase a MoviPrep Bowel Preparation Kit from a pharmacy. A prescription will be provided.

FOLLOW THE INSTRUCTIONS BELOW AND NOT THE INSTRUCTIONS THAT ARE INCLUDED WITH THIS KIT THE ENTIRE DAY BEFORE YOUR COLONOSCOPY – HAVE A CLEAR LIQUID DIET ONLY. NO SOLID FOODS.

Clear liquids include water, coffee, tea, Gatorade, carbonated and non-carbonated soft drinks. All the clear juices: apple, white grape. Jell-O, popsicles, and clear soups: bouillon or broth. Sugar or artificial sweeteners are allowed.

NOT ALLOWED ON A CLEAR LIQUID DIET: Dairy products: milk, cream, soy, artificial creamer, or pulpy juices like orange, grapefruit, pineapple, tomato and V-8. No alcoholic beverages.

STEP 1: AT 3:00 PM - MIX FIRST DOSE OF MoviPrep SOLUTION

- Empty **one** Pouch **A** and **one** Pouch **B** into the disposable container. **The other two packets are to be used for the second dose in Step 3 of this prep.**
- Add lukewarm drinking water to the top line of the container.
- Cap the bottle and shake to dissolve the powder.
- The mixed solution will be clear. Do not add anything other than the provided flavor packs to the solution.
- If you would like to drink the solution cold, store it in the refrigerator until it is time to start drinking.

STEP 2: AT 4:00 PM - START DRINKING THE MoviPrep SOLUTION:

- Drink one (8oz) glass every 30 minutes (about 4 glasses). Drink each glass steadily, at your own pace. **Do not “chug” it.** If you experience nausea, slow down in drinking the mix and try taking a few sips of one of the clear liquids listed above.
- A watery bowel movement should begin in approximately 1 hour or less. **Be sure to drink ALL the solution.**
- You may also have other clear liquids while drinking the MoviPrep solution.

STEP 3: At 6:00 PM - MIX THE REMAINING Pouch A and Pouch B following same instructions from STEP 1.

STEP 4: AT 7:00 PM - DRINK THE SECOND DOSE following the same instructions in **STEP 2.**



Garth LeCheminant, M.D.

David Morrow, M.D., F.A.C.S.

Iuliana Bobanga, M.D. F.A.C.S.

Jennifer Lemert, N.P.

2490 S. Woodworth Loop, Ste 450 Palmer, AK 99645

• Phone: (907) 745-8100 Fax: (907) 746-2655

• matusurgical.com

ON THE DAY OF YOUR COLONOSCOPY – You may continue to drink a small amount of clear liquids only up to **2 HOURS**
PRIOR TO ARRIVAL TIME then nothing by mouth.