



Garth LeCheminant, M.D. **David Morrow, M.D., F.A.C.S.** **Iuliana Bobanga, M.D., F.A.C.S.** **Jennifer Lemert, N.P.**
 2490 S. Woodworth Loop, Ste 450 Palmer, AK 99645 • Phone: (907) 745-8100 Fax: (907) 746-2655 • matsusurgical.com

PREPARATION INSTRUCTIONS FOR COLON SURGERY/with MoviPrep-Method

Someone from the hospital will call you **the day before** your surgery with the time to arrive.

_____ Mat-Su Regional Medical Center (907) 861-6858

PREPARATION INSTRUCTIONS FOR THE DAY BEFORE COLON-SURGERY

PLEASE READ THESE INSTRUCTIONS SEVERAL DAYS BEFORE YOUR PROCEDURE.

You will need the following items from a pharmacy: Prescriptions will be provided for a MoviPrep Bowel Preparation Kit and two antibiotics: Cipro & Flagyl.

FOLLOW OUR INSTRUCTIONS BELOW AND NOT THE INSTRUCTIONS THAT ARE INCLUDED WITH THIS KIT

THE ENTIRE DAY BEFORE YOUR COLON SURGERY – HAVE A CLEAR LIQUID DIET ONLY. NO SOLID FOODS.

Clear liquids include water, coffee, tea, Gatorade, carbonated and non-carbonated soft drinks. All the clear juices: apple, white grape. Jell-O, popsicles, and clear soups: bouillon or broth. Sugar or artificial sweeteners are allowed.

NOT ALLOWED ON A CLEAR LIQUID DIET: Dairy products: milk, cream, soy, artificial creamer, or pulpy juices like orange, grapefruit, pineapple, tomato and V-8. No alcoholic beverages.

STEP 1: AT 8:00 AM - MIX FIRST DOSE OF MoviPrep SOLUTION

- Empty **one** Pouch A and **one** Pouch B into the disposable container. **The other two packets are to be used for the second dose in Step 3 of this prep.**
- Add lukewarm drinking water to the top line of the container.
- Cap the bottle and shake to dissolve the powder.
- The mixed solution will be clear. Do not add anything other than the provided flavor packs to the solution.
- If you would like to drink the solution cold, store it in the refrigerator until it is time to start drinking.

STEP 2: AT 9:00 AM - START DRINKING THE MoviPrep SOLUTION:

- Drink one (8oz) glass every 30 minutes (about 4 glasses). Drink each glass steadily, at your own pace. **Do not “chug” it.** If you experience nausea, slow down in drinking the mix and try taking a few sips of one of the clear liquids listed above.
- A watery bowel movement should begin in approximately 1 hour or less. **Be sure to drink ALL the solution.**
- You may also have other clear liquids while drinking the MoviPrep solution.

STEP 3: AT 4:00 PM - MIX THE REMAINING Pouch A and Pouch B following same instructions from STEP 1.

STEP 4: AT 5:00 PM – Drink the second dose following the same instructions in STEP 2.

STEP 5. You will also start taking the antibiotics at the following times:

Cipro (ciprofloxacin) 500mg - 1 tablet at 11:00 am.

Flagyl (metronidazole) 500mg - 1 tablet at 11:00 am & 1 tablet at 9:00 pm.

ON THE DAY OF YOUR COLON SURGERY – You may continue to drink a small amount of clear liquids only up to **2 HOURS PRIOR TO ARRIVAL TIME** then nothing by mouth.