

3674 E. Meridian Loop Wasilla, AK 99654

907-373-8100 **** 907-373-3674 **=** matsusurgical.com **②**

- Garth LeCheminant, MD
- David Morrow, MD, FACS
- Iuliana Bobanga, MD, FACS
- Thomas Knolmayer, MD, FACS
- Jennifer Lemert, NP, CWOCN-AP

PREPARATION INSTRUCTIONS FOR COLON SURGERY/with SUPREP-Method

Someone from the hospital will call you the day before your surgery with the time to arrive.

Mat-Su Regional Medical Center 2500 S. Woodworth Loop Palmer, AK 99645 Phone: (907) 861-6338

PREPARATION INSTRUCTIONS FOR THE DAY BEFORE COLON-SURGERY

PLEASE READ THESE INSTRUCTIONS SEVERAL DAYS BEFORE YOUR PROCEDURE.

You will need the following items from a pharmacy: Prescriptions will be provided for a SUPREP Bowel Preparation Kit and two antibiotics: Cipro & Flagyl.

FOLLOW OUR INSTRUCTIONS BELOW AND **NOT** THE INSTRUCTIONS THAT ARE INCLUDED WITH THIS KIT

THE ENTIRE DAY BEFORE YOUR COLON SURGERY – HAVE A CLEAR LIQUID DIET ONLY. NO SOLID FOODS.

Clear liquids include water, coffee, tea, Gatorade, carbonated and non-carbonated soft drinks. All the clear juices: apple, white grape. Jell-O, popsicles, and clear soups: bouillon or broth. Sugar or artificial sweeteners are allowed.

NOT ALLOWED ON A CLEAR LIQUID DIET: Dairy products: milk, cream, soy, artificial creamer, or pulpy juices like orange, grapefruit, pineapple, tomato and V-8. No alcoholic beverages.

STEP 1: AT 9:00 AM – START DRINKING THE FIRST DOSE OF SUPREP SOLUTION

- Pour ONE (1) 6-ounce bottle of Suprep liquid into the 16-ounce cup
- Add cool drinking water to the 16-ounce line on the container and mix
- Drink ALL the liquid in the container
- IMPORTANT: You MUST drink two (2) more 16-ounce containers of water over the next hour.

STEP 2: AT 4:00 PM - START DRINKING THE SECOND DOSE SOLUTION

- You will drink your second dose of Suprep following the same instructions in STEP 1
- Drink ALL the liquid in the container
- IMPORTANT: You MUST drink two (2) more 16-ounce containers of water over the next hour.

STEP 3: You will also start taking the antibiotics at the following times:

Cipro (ciprofloxacin) 500mg – 1 tablet at 11:00 am.

Flagyl (metronidazole) 500mg – 1 tablet at 11:00 am & 1 tablet at 9:00 pm.

<u>ON THE DAY OF YOUR COLON SURGERY</u> – You may continue to drink a small amount of clear liquids only up to <u>2 HOURS</u> <u>PRIOR TO ARRIVAL TIME</u> then nothing by mouth.